

When can I stop?

You should keep taking your medication until your doctor tells you to stop.



You should keep taking it, even if you feel better.



If you want to stop taking it, talk to your doctor first.




Important

Some medications do not work well together.

You should always tell your doctor or pharmacist what you are taking.



If your medication looks different it might be a different make.



Alcohol can make you ill if you drink it when you are on medication.



Advice and Support

If you have any questions or worries about your medication talk to...

your doctor




or

the pharmacist at your local chemist



This leaflet was created by Michelle Forster, Betsy Wilkie, Dr Andre Strydom, Caroline Edwards and Dr Ian Hall.

The drawings are by Sally Poole, and from the CHANGE Picture Bank.




These people helped to make the leaflet easier to read: John Stevens, Jackie Ryan, Linda Doherty and Eamon Maher.



For more information contact The Elfrida Society, telephone: 020 7359 7443 fax: 020 7704 1358 Produced February 2001

Your medication is called

trifluoperazine or Stelazine



Typical antipsychotic

This is not an official manufacturer's patient information leaflet.

Why am I taking this?

It can help if you are anxious or paranoid.



It can help if you are too excited or 'hyper'.




It can stop you from hearing voices or seeing things that are not there.

How will it make me feel?

It should make you feel calm and sleep better.

It could take a month before you start to feel better.


What will it look like?



or




or



How do I take it?

Swallow your tablet or capsule whole with a glass of water.



Measure your liquid.



When do I take it?


Ask your doctor when to take it.

What if I forget to take it?

Take it as soon as you remember.



If it is nearly time for your next dose, only take one dose.




If you have any problems with taking your medication, you should talk to your doctor straight away.

Side effects

You might have some side effects, like...

- feeling sleepy
- a dry mouth
- too much spit in your mouth
- constipation
- put weight on
- your periods could change
- you could lose interest in sex



Remember, it is easy to get sunburnt when you are on trifluoperazine.

If you are worried, talk to your doctor.

IMPORTANT!

These side-effects are more serious...

- feeling faint
- things look funny
- a rash
- a fever
- stiffness
- shakiness
- problems with weeing

If you have any of these, you must talk to your doctor straight away!