
















Olanzapine

	<p>You are taking Olanzapine because you are hearing, seeing or sensing things that are not really there and stopping activities you normally do</p>
	<p>Olanzapine comes as tablets. Tablets should be taken with water</p>
	<p>When you take Olanzapine you may:</p> <ul style="list-style-type: none">Feel sleepyFeel hotFeel hungryFeel weakFeel stiff
	<p>See a doctor if you have a rash</p>
	<p>Olanzapine will help you feel calm and relaxed</p>






Pregablin

	<p>You are taking Pregablin because of feeling anxious or because you have epilepsy</p>
	<p>Pregablin come as tablets. Tablets should be taken with water</p>
	<p>When you take Pregablin you may:</p> <ul style="list-style-type: none"> Feel hungry Feel dizzy Put on weight Have a dry mouth
	<p>See a doctor if :</p> <ul style="list-style-type: none"> Your face or tongue swells up Your skin is red, blisters or peels
	<p>Pregablin will help you feel calm and relaxed</p>






Aripiprazol (Abilify)

	<p>You are taking Aripiprazol because you are hearing, seeing or sensing things that are not really there</p>
	<p>Aripiprazol comes as tablets. Tablets should be taken with water</p>
	<p>When you take Aripiprazol you may:</p> <ul style="list-style-type: none"> Feel dizzy or weak Feel restless Feel sick Have problems sleeping Have a headache Become constipated
	<p>See a doctor if you:</p> <ul style="list-style-type: none"> feel stiff feel shaky have blurred vision have a racing heart
	<p>Aripiprazol will help you feel calm and relaxed</p>






Celebrex

	<p>You are taking Celebrex because you are in pain</p>
	<p>Celebrex comes as tablets. Tablets should be taken with water.</p>
	<p>When you take Celebrex you may: have a sore throat have tummy pains feel sick have problems sleeping</p>
	<p>See a doctor if: you have problems breathing your face swells up you have a rash</p>
	<p>Celebrex will help to ease the pain</p>






Lorazepam

	<p>You are taking Lorazepam because you are feeling anxious</p>
	<p>Lorazepam comes as tablets. Tablets should be taken with water.</p>
	<p>When you take Lorazepam you may:</p> <ul style="list-style-type: none">Feel unsteadyFeel light-headed and faintFeel sleepy and slowFeel confusedHave a headache
	<p>See a doctor if you:</p> <ul style="list-style-type: none">Have a rash or blotches on your skin
	<p>Lorazepam will help you to feel calm and relaxed</p>






Procyclidine

	<p>You are taking Procyclidine because your other medications are making you feel stiff or shaky</p>
	<p>Procyclidine comes as tablets or liquid. Tablets should be taken with water</p>
	<p>When you take Procyclidine you may:</p> <ul style="list-style-type: none">Have blurred visionHave a dry mouthHave problems to weeFeel sick
	<p>See a doctor if you:</p> <ul style="list-style-type: none">Have unusual body movementsHave uncontrolled movements of the face and tongue
	<p>Procyclidine will stop you feeling stiff or shaky</p>






Lithium

	<p>You are taking Lithium because you are feeling unwell</p>
	<p>Lithium comes as tablets or liquid. Tablets should be taken with water</p>
	<p>When you take Lithium you may:</p> <ul style="list-style-type: none"> Need to drink more Need to wee more Feel confused Feel sick Feel sleepy Feel unsteady on your feet
	<p>See a doctor if you:</p> <ul style="list-style-type: none"> Have dizziness or blurred vision Hear ringing in your ears Feel weak or numb in your arms and legs Have sickness and diarrhoea
	<p>Lithium will help you to feel better</p>






Fluoxetine

	You are taking Fluoxetine because you have depression or obsessive-compulsive disorder
	Fluoxetine comes as tablets or liquid. Tablets should be taken with water
	When you take Fluoxetine you may: Have problems sleeping Have an upset tummy Feel sick Feel confused Need to wee more
	See a doctor if you have: A rash all over your skin Itchy skin A fever
	Fluoxetine will help you feel calm and relaxed

Diazepam

	<p>You are taking Diazepam because of feeling anxious or you have epilepsy</p>
	<p>Diazepam comes as tablets and liquid. Tablets should be taken with water</p>
	<p>When you take Diazepam you may:</p> <ul style="list-style-type: none">Feel unsteadyFeel light-headed and faintFeel sleepy and slowFeel confusedHave a headache
	<p>See a doctor if you:</p> <ul style="list-style-type: none">Have a rash or blotches on your skin
	<p>Diazepam will help you to feel calm and relaxed</p>

Risperidone

	<p>You are taking Risperidone Because you have a condition that affects the way you think, feel and act</p>
	<p>Risperidone comes as tablets or liquid. Tablets should be taken with water</p>
	<p>When you take Risperidone you may: Have a dry mouth Feel drowsy Feel restless Have problems sleeping</p>
	<p>See a doctor if you: Feel faint Have a rash Feel stiff or shaky Have a pounding heart</p>
	<p>Risperidone will help you feel calm and relaxed</p>