













Clopixol

	<p>You are taking Clopixol due to feeling unwell</p>
	<p>Clopixol comes as an injection, it will be injected into your bottom</p>
	<p>When you take Clopixol you may: Feel stiff Gain weight Become confused Have blurred vision</p>
	<p>See a doctor if you: Have difficulties breathing Itchy skin Swelling of the lips, face or tongue</p>
	<p>Clopixol will help you feel better</p>






Haloperidol

	<p>You are taking Haloperidol because you have a condition that affects the way you think, feel and do things</p>
	<p>Haloperidol comes as an injection and will be injected into your bottom</p>
	<p>When you take Haloperidol you may:</p> <ul style="list-style-type: none">Have a rashFeel sickFind it hard to weeHave a dry mouth
	<p>See a doctor if:</p> <ul style="list-style-type: none">You have swelling, pain and redness in the legsSwelling of the face or throatJerky movements
	<p>Haloperidol will help you feel better</p>






Movicol

	<p>You are taking Movicol because you suffer from constipation.</p>
	<p>Movicol comes in a powder that is mixed with water.</p>
	<p>Movicol may make you: Feel sick Have tummy pains Have a sore bottom</p>
	<p>See a doctor if: Have an itchy skin Have problems breathing</p>
	<p>Movicol will help you to be able to go to toilet.</p>






Carbamazepine

	<p>You are taking Carbamazepine because you have epilepsy or a mood disorder</p>
	<p>Carbamazepine comes as tablets or liquid. Tablets should be swallowed with a glass of water</p>
	<p>When taking Carbamazepine you may:</p> <ul style="list-style-type: none">Feel sickHave tummy painPut weight onFeel unsteady
	<p>See a doctor if:</p> <ul style="list-style-type: none">You have swollen anklesYou have difficulty breathingYour face swells upYou have a rash
	<p>Carbamazepine will help you to feel better</p>






Omeprazole

	<p>You are taking Omeprazole because you have indigestion or heart burn</p>
	<p>Omeprazole comes as a tablet. Tablets should be swallowed with a glass of water</p>
	<p>While taking Omeprazole you may:</p> <ul style="list-style-type: none">Feel sickHave a headacheFeel tiredHave tummy pains
	<p>See a doctor if :</p> <ul style="list-style-type: none">You have swelling of the lips, face or throatYou have a feverYour skin blisters
	<p>Omeprazole will help to feel more comfortable</p>






Sodium valporate

	<p>You are taking Sodium Valporate because you have epilepsy</p>
	<p>Sodium Valporate comes as a liquid, tablets or as granules. The tablets should be taken with a glass of water the granules should be sprinkled onto a small amount of soft food or into a cold drink.</p>
	<p>While taking Sodium Valporate you may:</p> <ul style="list-style-type: none">Feel sickHave tummy painsPut on weightFeel faint
	<p>See a doctor if:</p> <ul style="list-style-type: none">You have a rashYou face, lips, tongue or throat swell upYou have problems breathing
	<p>Sodium Valporate will help to reduce the number of epileptic fits you have</p>






Ibuprofen

	<p>You are taking Ibuprofen because you are in pain</p>
	<p>Ibuprofen comes as tablets. Tablets should be taken with water after food</p>
	<p>While taking Ibuprofen you may: Have a rash Have tummy pains Feel sick Become constipated</p>
	<p>See a doctor if: You have difficulty breathing Your face tongue or throat start to swell up</p>
	<p>Ibuprofen will help the pain get better</p>






Clomipramine

	<p>You are taking Clomipramine because you have depression, or a phobia</p>
	<p>Clomipramine comes as tablets. Tablets should be taken with water</p>
	<p>While taking Clomipramine you may: Feel hungry Put weight on Feel sick Have problems to wee</p>
	<p>See a doctor if: You have a face You have difficulty breathing</p>
	<p>Clomipramine will help you to feel better</p>






Docosate Sodium

	<p>You are taking Docosate Sodium because you are constipated or to stop you becoming constipated</p>
	<p>Docosate Sodium comes as a tablet. Tablets should be swallowed whole with a glass of water</p>
	<p>While taking Docosate Sodium you may:</p> <ul style="list-style-type: none">Feel sickHave tummy painsHave a rash
	<p>See a doctor if any of the side effect last for more then a few days or become a problem</p>
	<p>Docosate Sodium will help you to go to the toilet</p>






Paracetamol

	<p>You are taking Paracetamol because you are in pain</p>
	<p>Paracetamol comes as tablets and liquid. Tablets should be swallowed with a glass of water</p>
	<p>When taking Paracetamol you may: Have a rash Have bleeding gums</p>
	<p>See a doctor if: You have difficulty breathing The rash gets worse Your face tongue or throat swells up</p>
	<p>Paracetamol will help the pain get better</p>




Melatonin

	<p>You are taking Melatonin because you have problems getting to sleep or staying asleep</p>
	<p>Melatonin comes as tablets and should be taken with food 1-2 hours before you go to bed. Tablets should be taken whole with a glass of water</p>
	<p>While taking Melatonin you may: Become constipated Have tummy pains Feel sick</p>
	<p>See a doctor if the symptoms get worse</p>
	<p>Melatonin will help you to sleep</p>

Senna

	<p>You are taking Senna because you are constipated</p>
	<p>Senna comes as tablets or as syrup. Tablets should be taken with water</p>
	<p>While taking Senna you may: Lose weight Have diarrhoea Have constipation</p>
	<p>See a doctor if any of the symptoms get worse</p>
	<p>Senna will help you to be able to go to the toilet</p>

Lactulose

	<p>You are taking Lactulose because you are constipated</p>
	<p>Lactulose is a liquid</p>
	<p>While taking Lactulose you may: Have diarrhoea Feel bloated</p>
	<p>See a doctor if the side effects get worse</p>
	<p>Lactulose will help you to go to toilet</p>